

Maximum Strength Mucinex® Fast-Max® NIGHT TIME Cold & Flu Liquid

Drug Facts

Active ingredients (in each 20 mL) Purposes

Acetaminophen 650 mg... Pain reliever/fever reducer
 Diphenhydramine HCl 25 mg
 Antihistamine/cough suppressant
 Phenylephrine HCl 10 mg..... Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough ■ nasal congestion
 - minor aches and pains ■ sore throat
 - headache ■ runny nose ■ sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease ■ heart disease
- high blood pressure ■ thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- breathing problems such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

Drug Facts (continued)

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- tamper evident: do not use if neckband on bottle cap is broken or missing
- store between 20-25°C (68-77°F)
- do not refrigerate ■ dosing cup provided

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
 *may contain this ingredient

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.