

Maximum Strength Mucinex® Fast-Max® NIGHT TIME Cold & Flu Caplets

Drug Facts

Active ingredients (in each caplet) Purposes

Acetaminophen 325 mg ... Pain reliever/fever reducer
Diphenhydramine HCl 25 mg Antihistamine
Phenylephrine HCl 5 mg Nasal decongestant

Uses ■ temporarily relieves these common cold and flu symptoms: ■ nasal congestion
■ headache ■ minor aches and pains
■ sore throat ■ runny nose and sneezing
■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 12 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks daily while using this product

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
 - pain or nasal congestion gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present ■ new symptoms occur
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see Overdose warning)

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients corn starch, croscarmellose sodium, crospovidone, FD&C Blue #1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, triacetin

Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.