

# Maximum Strength Mucinex® Fast-Max®

## DAY TIME Severe Cold and

## NIGHT TIME Cold & Flu Liquids

### Drug Facts

#### Active ingredients (in each 20 mL) Purposes Mucinex FAST-MAX Day Time Severe Cold

Acetaminophen 650 mg ..... Pain reliever/fever reducer  
 Dextromethorphan HBr 20 mg ..... Cough suppressant  
 Guaifenesin 400 mg ..... Expectorant  
 Phenylephrine HCl 10 mg ..... Nasal decongestant

#### Active ingredients (in each 20 mL) Purposes Mucinex FAST-MAX Night Time Cold & Flu

Acetaminophen 650 mg ..... Pain reliever/fever reducer  
 Diphenhydramine HCl 25 mg ..... Antihistamine/  
 cough suppressant  
 Phenylephrine HCl 10 mg ..... Nasal decongestant

### Uses

- temporarily relieves these common cold and flu symptoms:
  - nasal congestion
  - cough
  - minor aches and pains
  - headache
  - sore throat
- runny nose and sneezing (*Night Time Cold & Flu only*)
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (*Day Time Severe Cold only*)
- temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs that contain acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin (*Night Time Cold & Flu only*)
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

#### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma (*Night Time Cold & Flu only*)
- a breathing problem such as emphysema or chronic bronchitis (*Night Time Cold & Flu only*)

### Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (*Night Time Cold & Flu only*)

**When using this product** ■ **do not use more than directed**  
 ■ excitability may occur, especially in children (*Night Time Cold & Flu only*)

- marked drowsiness may occur (*Night Time Cold & Flu only*)
- alcohol, sedatives, and tranquilizers may increase drowsiness (*Night Time Cold & Flu only*)
- avoid alcoholic drinks (*Night Time Cold & Flu only*)
- be careful when driving a motor vehicle or operating machinery (*Night Time Cold & Flu only*)

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided with product
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

### Other information

- dosing cup provided
- each 20 mL contains: **sodium 12 mg**
- store between 20-25°C (68-77°F) ■ do not refrigerate

#### Inactive ingredients (Mucinex FAST-MAX Day Time Severe Cold)

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate\*, xanthan gum \*may contain this ingredient

#### Inactive ingredients (Mucinex FAST-MAX Night Time Cold & Flu only)

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate\*, xanthan gum \*may contain this ingredient