

Maximum Strength Mucinex® Fast Max® Severe Congestion & Cough/ Night Time Cold & Flu Caplets

Drug Facts

Active ingredients (in each caplet) Purposes Mucinex FAST-MAX DAY TIME

Severe Congestion & Cough

Dextromethorphan HBr 10 mg..... Cough suppressant
Guaifenesin 200 mg..... Expectorant
Phenylephrine HCl 5 mg..... Nasal decongestant

Active ingredients (in each caplet) Purposes Mucinex FAST-MAX NIGHT TIME Cold & Flu

Acetaminophen 325 mg..... Pain reliever/fever reducer
Diphenhydramine HCl 25 mg..... Antihistamine
Phenylephrine HCl 5 mg..... Nasal decongestant

Uses

- temporarily relieves (**DAY TIME only**):
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep
 - nasal congestion due to a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (**DAY TIME only**)
- temporarily relieves these common cold and flu symptoms (**NIGHT TIME only**):
 - nasal congestion
 - minor aches and pains
 - headache
 - sore throat
 - runny nose and sneezing
- temporarily reduces fever (**NIGHT TIME only**)
- controls cough to help you get to sleep

Warnings

- Liver warning (NIGHT TIME only):** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 12 caplets in 24 hours, which is the maximum daily amount
 - with other drugs that contain acetaminophen
 - 3 or more alcoholic drinks daily while using this product

Sore throat warning (NIGHT TIME only): If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. (**NIGHT TIME only**)
- with any other drug containing diphenhydramine, even one used on the skin (**NIGHT TIME only**)
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease (**NIGHT TIME only**)
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma (**NIGHT TIME only**)
- a breathing problem such as emphysema or chronic bronchitis (**NIGHT TIME only**)

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema (**DAY TIME only**)
- cough that occurs with too much phlegm (mucus) (**DAY TIME only**)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin (**NIGHT TIME only**)
- taking sedatives or tranquilizers (**NIGHT TIME only**)

When using this product do not use more than directed.

In addition, for **NIGHT TIME only**:

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with a fever (**DAY TIME only**)
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days (**NIGHT TIME only**)
- fever gets worse, or lasts more than 3 days (**NIGHT TIME only**)
- redness or swelling is present (**NIGHT TIME only**)
- new symptoms occur (**NIGHT TIME only**)
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. (**NIGHT TIME only**) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see **Overdose warning**)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients (Mucinex FAST-MAX DAY TIME Severe Congestion & Cough)

croscarmellose sodium, FD&C Blue #2 aluminum lake, FD&C Red #40 aluminum lake, methacrylic acid-ethyl acrylate copolymer, mica, microcrystalline cellulose, polyethylene glycol 3350, polysorbate 80, polyvinyl alcohol, povidone K29/32, sodium bicarbonate, talc, titanium dioxide

Inactive ingredients (Mucinex FAST-MAX NIGHT TIME Cold & Flu)

corn starch, croscarmellose sodium, crospovidone, FD&C Blue #1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, triacetin