

# Maximum Strength Mucinex® Sinus-Max™ Pressure & Pain Caplets

## Drug Facts

### Active ingredients Purpose (in each caplet)

Acetaminophen 325 mg	Pain reliever
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

### Uses

- temporarily relieves:
  - nasal congestion
  - headache
  - minor aches and pains
  - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

## Drug Facts (continued)

- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not use more than directed**

### Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

■ do not take more than directed (see Overdose warning)

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

### Other information

- store at 20-25°C (68-77°F)

**Inactive ingredients** croscarmellose sodium, crospovidone, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

### Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.