

Maximum Strength Mucinex[®] Sinus-Max[®] Pressure & Pain Liquid

Drug Facts

Active ingredients (in each 20 mL)	Purposes
---	-----------------

Acetaminophen 650 mg	Pain reliever
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- Temporarily relieves:
 - nasal congestion ■ headache
 - minor aches and pains
 - sinus congestion and pressure
 - temporarily promotes nasal and/or sinus drainage
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease

- heart disease ■ high blood pressure
- thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus) ►

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- adults and children 12 years and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: **sodium 12 mg**
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum
*may contain this ingredient