

# Children's MUCINEX<sup>®</sup>

## DAY TIME MULTI-SYMPTOM COLD/ NIGHT TIME MULTI-SYMPTOM COLD Liquids

### Day Time Multi-Symptom Cold – Age 4+

#### Drug Facts

##### Active ingredients (in each 5 mL) Purposes

Dextromethorphan HBr 5 mg.....Cough suppressant  
Guaifenesin 100 mg.....Expectorant  
Phenylephrine HCl 2.5 mg.....Nasal decongestant

#### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help your child get to sleep
  - nasal congestion due to a cold
  - stuffy nose

#### Warnings

**Do not use** in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

##### Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

**When using this product** do not use more than directed

##### Stop use and ask a doctor if

- your child gets nervous, dizzy or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

#### Other information

- each 5 mL contains: **sodium 3 mg**
- tamper evident: do not use if neckband on bottle cap is broken or missing
- store between 20-25°C (68-77°F)
- do not refrigerate ■ dosing cup provided ▶

### Day Time Multi-Symptom Cold – Age 4+ (cont'd)

#### Drug Facts (continued)

**Inactive ingredients** anhydrous citric acid, edetate disodium, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate\*, xanthan gum  
\*may contain this ingredient

#### Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

### Night Time Multi-Symptom Cold – Age 6+

#### Drug Facts

##### Active ingredients (in each 10 mL) Purposes

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Diphenhydramine HCl 12.5 mg.....Antihistamine/  
cough suppressant  
Phenylephrine HCl 5 mg.....Nasal decongestant

#### Uses

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - headache ■ sore throat
  - nasal congestion ■ runny nose
  - sneezing ■ cough
- controls cough to help your child get to sleep
- temporarily reduces fever

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
  - with other drugs containing acetaminophen
- Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

##### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- with any other drug containing diphenhydramine, even one used on the skin
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

##### Ask a doctor before use if your child has

- liver disease ■ heart disease
- diabetes ■ high blood pressure
- thyroid disease ■ glaucoma ▶

### Night Time Multi-Symptom Cold – Age 6+ (cont'd)

#### Drug Facts (continued)

- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

##### Ask a doctor or pharmacist before use if your child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

##### When using this product

- do not use more than directed (see **Overdose warning**)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

##### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

##### Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see Overdose warning)**
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

#### Other information

- each 10 mL contains: **sodium 6 mg**
- store between 20-25°C (68-77°F)
- dosing cup provided
- do not refrigerate

**Inactive ingredients** anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate\*, xanthan gum \*may contain this ingredient

#### Questions? 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.